


Physical Health Assessment and Monitoring for COVID-19



A Guide for Nurses in Community Mental Health Settings

Contents

Acknowledgements

A special thank you to Catherine Gamble who is leading the development of the 5 & 1 V mental health nursing COVID-19 guidance and information. RCN Representative Stephen Jones for creating and coordinating this guidance. Geoff Brennan for his comprehensive contribution to the overall content.

Catherine Gamble
Professional Lead for Mental
Health,
Royal College of Nursing

Stephen Jones
Project Development Nurse,
South London Mental Health &
Community Partnership

Geoff Brennan
Chief Executive, BRIGHT
Executive Director, STAR WARDS

The RCN would like to thank everyone who provided their expertise and guidance to develop the information set out in this document.

Tim Coupland
Clinical Rep for Parity of Esteem,
Royal College of Nursing

Hannah MacDonald
Lived Experience Representative,
West London NHS Trust

Prof David Evans , OBE, NTF
Professor in Sexualities and
Genders: Health & Wellbeing,
University of Greenwich

Yaw Mensah
Mental Health Nurse,
Oxleas NHS Foundation Trust

Dr Dawne Garrett
Professional Lead Care of Older
People and Dementia Care,
Royal College of Nursing

Dr Ehab Morgan
Consultant Psychiatrist,
Oxleas NHS Foundation Trust

Matthew WarET Q q 49.4 85.5NHS Fc

Mary Jones
Senior Community Learning
Disability Nurse,
Kent Community Health NHS
Foundation Trust

Introduction

The Coronavirus pandemic presents health care services with unique dilemmas. These are unprecedented times for which there is no clear rulebook. Our guidance, therefore, can only

Shielding: High-Risk and Extremely Vulnerable

The UK government has provided [guidance on shielding and protecting people defined on medical grounds as extremely vulnerable from COVID-19](#) (March, 2020). The guidance has defined people as high-risk of severe illness from coronavirus (COVID-19) because of an underlying health condition if

People falling into this high-risk and extremely vulnerable group include:

1. Solid organ transplant recipients.
2. People with specific cancers:
 - people with cancer who are undergoing active chemotherapy or radical radiotherapy for lung cancer
 - people with cancers of the blood or bone marrow such as leukaemia, lymphoma or myeloma who are at any stage of treatment
 - people having immunotherapy or other continuing antibody treatments for cancer
 - people having other targeted cancer treatments which can affect the immune system, such as protein kinase inhibitors or PARP inhibitors
 - people who have had bone marrow or stem cell transplants in the last 6 months, or who are still taking immunosuppression drugs
3. People with severe respiratory conditions including all cystic fibrosis, severe asthma and severe COPD.
4. People with rare diseases and inborn errors of metabolism that significantly increase the risk of infections (such as SCID, homozygous sickle cell).
5. People on immunosuppression therapies sufficient to significantly increase risk of infection.
6. Women who are pregnant with significant heart disease, congenital or acquired.

The NHS also includes conditions that make people more likely to get infections (i.e. diabetes and HIV), as well as people who are taking medication that weakens the immune system (i.e. Clozapine and Lithium).


Supporting Community Services

With the Royal Collage of Psychiatrists, RCN members have formulated a [clinical decision framework](#) to enable the allocation of care and treatment for community mental health patients during the COVID-19 outbreak. We have also produced an [algorithm](#) to support clinical decision-making for service users with high-risk physical health needs in the community.

High-risk needs are present in people who are either considered extremely vulnerable and/or are prescribed high-risk medication (i.e. Clozapine & Lithium). Those considered to be at very high-risk fit into the high-risk categories; in addition, live alone and are unable to monitor their own physical health needs (see figure 1 on page 6).

Guidance for DVVHVVLQJ D SHUVRQ V high-risk Very DO KHDOWK ULVN

Algorithm: Supporting Clinical Decision -Making amid COVID-19



Appendices

UsefulResources

RCN Guidance - COVID-19: <https://www.rcn.org.uk/clinical-topics/mental-health/covid-19-guidance>

RCN Guidance for Mental Health Nurses - COVID-19: <https://www.rcn.org.uk/clinical-topics/mental-health/covid-19-guidance>

RCPsych COVID-19 guidance for mental health clinicians:

<https://www.rcpsych.ac.uk/about-us/responding-to-covid-19/responding-to-covid-19-guidance-for->

The RCN represents nurses and nursing, promotes excellence in practice and shapes health policies

RCN Online

www.rcn.org.uk

RCN Direct

www.rcn.org.uk/direct

0345 772 6100

Published by the Royal College of Nursing

20 Cavendish Square London
W1G 0RN

020 7409 3333

RCN Legal Disclaimer

This publication contains information, advice and guidance to help members of the RCN. It is intended for use within the UK but readers are advised that practices may vary in each country and outside the UK. The information in this booklet has been compiled from a range of sources and is not intended to be a substitute for professional advice.