y C g

1 .

Abthisdcen

- hysteroscopy
- nurses working within early pregnancy and acute gynaecology settings
- and as part of any extended role in history taking and examination for the assessment of symptomatic women.

Training is intended to enable registered health care professionals to demonstrate competence at recognising the abnormal or symptomatic from the normal or

removal of a foreign body, vaginal dilatation, fitting of contraceptive devices, removal of placenta, evacuation of retained products, transvaginal chorionic villus sampling, endometrial ablation, and assisted reproduction techniques such as insemination or embryo transfer.

BACK TO CONTENTS

Registered NAs must meet the core clinical competencies in the Skills for Health competency framework set out in the cervical screening sample taker guidance at: gov.uk/government/publications/cervical-screening-cervical-sample-taker-training .

To undertake cervical screening, NAs must have:

- completed a nursing associate qualification and be registered as a NA with the Nursing and Midwifery Council (NMC)
- undertaken initial theory and practical training as required by the NHS CSP, successfully completed the course and assessed as competent
- undertaken update training and maintained competency in line with national cervical sample taker training guidance.

Local governance: The registered NA role is not yet a named profession under the Treatment of Disease, Disorder or Injury (TDDI) legislation regulated by the Care Quality Commission (CQC). However, the CQC expects any provider to consider safety, quality, competency and TDDI legislation when deploying a NA. See: cqc.org.uk/sites/default/files/20190123_briefing_for_providers_nursing_associates_0.pdf CQC briefing for providers.

When a NA has registered with the NMC, a registered professional listed under the legislation (registered nurse or GP) will need to supervise the practice of that NA. The supervisor must be present at the GP practice when the NA is carrying out the procedure. The supervisor can undertake indirect supervision of the NA when carrying out this procedure. This is a delegated activity and the NA would be expected to work within the remits of their professional code.

NHS England/Improvement, Health Education England and PHE are working together to follow a test cohort of registered nursing associateen-GBrnursisisp, 6 (s a 9 (r)-9p)-2.8 (n)-0.9 (16.9 (t).1 (e)

NHS 2.7 (n)5.8 (g)5.3 ()-s'st.1 (h)-1.166 (e i)6.9 s-1 (e N)4.82.2 (H)red nuretemland aner to

When the chaperone is a nurse or another member of the health care team, they can act as an advocate for the woman to:

- explain what will happen during the examination or procedure, and the reasons why
- assess the woman's understanding of what she has been told
- provide a reassuring presence during the examination or procedure
- safeguard against any pain, humiliation, intimidation or unnecessary discomfort
- observe the woman to ensure she feels safe and is comfortable
- always be sensitive to cultural and religious issues, as well as the context of the woman's circumstances and specific needs.

Although a woman should be offered a chaperone, she may find it difficult to have a

BACK TO CONTENTS

- How will the information obtained be used to benefit the woman?
- Are you competent to perform the planned procedure?
-

- explain that she may stop the examination at any point with a request to do so, and agree how that request can be made, such as a key word, or raising of the hand
- it is recommended that a woman should be advised how to take her own low-vaginal swab for chlamydia and gonorrhoe (this is common practice in sexual health services) BEFORE the rest of the examination
- explain that some women may experience some spotting after swabs and cervical sampling, if appropriate
- ensure the woman has privacy if she needs to undress and show her where to place her clothing
- the woman should be advised that it is usually only necessary to remove her lower garments
- assistance to remove garments should only be given if required, and not in an attempt to hurry the woman
- ensure the woman has enough tissue or a sheet to cover the genital area when undressed
- ask if she would like you to talk through the procedure as it happens
- ask the woman to let you know when she is ready.

If you are preparing the area, make sure the woman is aware of the possible sounds she might hear and what they represent:

- position and check trolley, and the availability of required equipment
- ensure there is good light and that any viewing light is switched on
- light sources should be cold light and should not have hot exteriors which may cause discomfort to the woman
- wash hands and wear gloves – consider possible latex allergy.

The examination

The following is broad guidance for vaginal and genital examination. The examinations performed will differ between fields of practice, die lr dig ihe wre losif phe wpf. 8.2 ((. 8.2i((2)-3.8 ((e)3.2 ((e)-

umbilicus towards pubic bone to identify the uterus, and note findings (size, position and tenderness of uterus). For the midwife, this examination will also focus on identifying the lie, position and gestation of a pregnancy.

- Palpate the abdomen in a systematic manner.
- Gloves are not required for this part of the examination.
- Light palpation – feeling the abdomen gently is especially helpful in identifying tenderness and muscular resistance. Keep your hand and forearm on a horizontal plane, with fingers together and flat on the abdominal surface, palpate the abdomen with a light, gentle dipping motion. When moving your hand from place to place, raise it off the skin. Moving smoothly, feel in all quadrants.
- Deep palpation – if there is no evidence of distress during light palpation, palpate the abdomen in the same systematic manner but deeper.
-

This examination is used by appropriately trained nurses and midwives, mainly for assessment and diagnostic purposes.

- Insert a gloved and lubricated index and/or middle finger into the vagina (depending on the vaginal canal may only be able to insert one digit).
- Assess the vagina and note findings such as vaginal tone and vaginal wall support – degree of prolapse, varicosities, tenderness, protrusions, foreign bodies etc.
- Place the other hand on the abdomen and press towards the fingers inside the vagina.

Examination of the cervix:

- locate the cervix and lightly feel this between two fingers, then assess its size and movement; it should move freely (if there is an infection present and/or cervical motion tenderness is a sign of Pelvic Inflammatory Disease (PID) and would need further assessment by a doctor immediately
- palpate the cervix – it should feel smooth and firm (hard and lateral displacement could indicate the presence of tumours/fibroids)
- comment on findings such as mobility, discomfort, size, shape, consistency, depth/ projection into vagina, angle, or any masses.

Palpation of the uterus:

- the abdominal hand should be midway between the umbilicus and the symphysis pubis
- the vaginal hand with palmar surface facing anteriorly should maintain contact with the cervix so that the nurse/midwife examining the woman can use the cervix as a ‘landmark’ for palpating the uterus
- lightly applying pressure to the posterior portion of the cervix with the vaginal hand will bring the uterus towards the abdomen
- once the uterus is raised, use the external hand to palpate, taking note of size, shape, position and consistency
- if the uterus is retroverted or retroflexed, it will not come up between the examining hands – differentiation between an anteverted and retroverted uterus is vital for certain procedures
- assess the uterus, taking note of size, shape, position, consistency, mobility and tenderness
- record findings from the uterine palpation.

Follow the guidelines below for palpation and examination of the adnexa:

- move abdominal hand to the lower abdominal quadrant on the same side as the internal hand
- move fingers in the vagina to either the right or left sides of the lateral fornix
- apply firm and steady pressure, beginning medial to the anterior iliac crest
- note any tenderness or masses

BACK TO CONTENTS

- palpate the position of the fallopian tubes on either side of the uterus; these are not normally palpable or tender
- when palpating the ovaries at the end of the fallopian tubes, advise the woman that some discomfort is likely
- the ovaries are approximately 2-4 cm in length, smooth, firm, mobile, sensitive to touch but not tender and, if palpable, should feel the size of an almond. In post-menopausal women they are smaller
- gentle moving of the cervix slightly from side to side will demonstrate ‘cervical excitation’. Should there be any adnexal masses or tenderness, advice should be sought
- if there is a potential diagnosis of ectopic pregnancy, this procedure should be performed by a skilled clinician, following pregnancy testing and ultrasound scanning and requires immediate referral to a specialist if there is any suspicion of an ectopic pregnancy.

Fllig the exiatn

Switch off the examination light and provide privacy for the woman to get dressed or rearrange her clothing. Ensure the woman has tissue available to wipe away any lubricant or discharge and that there is access to washing facilities and sanitary pads, if needed.

Ensure a full record is made of the examination performed, and that any tests taken and findings observed are recorded clearly and contemporaneously in the woman’s notes. Wash hands and document your findings using local policy. The following points should be included in your records:

- abdomen
- external genitalia
- vagina
- cervix
- uterus
- adnexae.

And may include reference to:

- size
- positi58Tw 11 0 0 11 70.8661 256.6681 Tm (•)Tj ET BT /Span <</ActualText (þ024 Tw 11 0 0 11 85.0394 175.

BACK TO CONTENTS

Some women will have a history of traumatic experiences with previous examinations or may have experienced sexual abuse, physical abuse or rape in the past. This may be

see the RCN publication Female Genital Mutilation: An RCN resource for nursing and midwifery practice (RCN, 2019).

FGM is illegal in the UK, and widely recognised as a form of abuse. The legislation relating to FGM has changed in 2015, and all health care practitioners should be aware of their role and responsibility with regard to reporting and recording, as well as how to best care for any girl or woman affected by FGM.

If a girl under 18 years of age has or is suspected of having had FGM carried out, then local safeguarding procedures should be implemented, including informing the police via the 101 nonemergency number (DH, 2015).

There is now a mandatory duty for all regulated health care professionals to report any concerns they may have about a female under 18 years and record when FGM is disclosed or identified as part of NHS health care.

If over 18 years of age recording of the case should be carried out in line with Department of Health requirement under the Enhanced Dataset requirement (HSCIC, 2018). Further information is availably from the Department of Health (DH, 2019). The priority for the woman should always be proving the best care possible to support any physical or psychological, or psycho sexual needs she may have (RCN, 2019).

Virginity testing and hymenoplasty have been illegal in the UK since 2022 (Sections 136-160 of the Health and Care Act (on virginity testing and hymenoplasty 1 July 2022 (Gov.uk, 2022a)). This legislation was introduced following growing concern that some girls and young women are being forced/coerced to have a virginity test, and subsequently undergoing hymen repair surgery.

Nurses need to understand the implications of this change in legislation, which may impact on their practice.

The government published guidelines in July 2022, which can be found at: www.gov.uk/government/publications/virginity-testing-and-hymenoplasty-multi-agency-guidance (Gov.uk, 2022b).

Virginity testing , also referred to as hymen, ‘two-finger’ or vaginal examination, is an inspection of the female genitalia, intended to determine whether a woman or girl has had vaginal intercourse.

Hymenoplasty (also known as hymenorrhaphy, hymen reconstruction or hymen repair) is a surgical intervention that involves reconstructing the hymen. There are a number of different techniques to achieve this but generally it involves stitching the torn edges of the hymen together with dissolvable stitches. The aim of the procedure is to ensure that the woman bleeds when she next has sexual intercourse (DHSE, 2021).

Hymenoplasty is regarded as a form of cosmetic/plastic surgery and used in the same way as other forms of cosmetic surgery for aesthetic rather than medical reasons, hence part of the controversy about banning it rests with the issue of freedom to choose by adult women (and those who were born female, but may not identify as being female), in the same way that there is freedom to choose other forms of surgery for aesthetic reason.

This is a contentious issue but should a woman be restrained, for example, in a custodial setting, then the nurse is still responsible for ensuring consent is given for the procedure to be carried out and that the woman's dignity is maintained.

Some women will request to only be examined by a female and this should be respected. If a female health professional has been requested but is unavailable, alternative arrangements may have to be made. In emergency situations, where no female health professionals are available, sensible and practicable measures must be taken.

If the woman gains sexual satisfaction from the examination the need for a chaperone is paramount. This should be clearly documented.

If a pregnant woman has had an antepartum haemorrhage or is known to have placenta praevia then an examination should not be carried out.

Women with a limited command of or no English will require a recognised translator. Due to the intimacy of the examination and the sensitivity of the consultation, a family member or friend may not be appropriate to assist with the translation The general advice is that family members should not be used for translation purposes. If translated forms of written information are available they should be provided prior to the examination.

A woman should give consent to "examination under anaesthesia" and be made aware of, as well as have the right to refuse any teaching or training of medical, nursing or midwifery learners whilst anaesthetised. Women can feel particularly vulnerable about being under anaesthetic and not having any control over the situation. It is therefore necessary that nothing additional is performed other than what is consented for. The be11.7 (or)36.8 (n)40.9 (n 8.179i)15 0.013 Tw 12 0 0 32exan14.17 (t 93(s)7.pan (e)-0.8 72.1 (e11.7 (or)369r)

BACK TO CONTENTS

3. [illegible]

[illegible]

5. Demonstrate the knowledge and skills required to safely and effectively perform genital bimanual examination.
6. D

4. Step

electronic feedback mechanisms such as an online survey tool. The learner also should complete an evaluation of their training and return to the assessor for feedback on the process (Appendix 5).

The logbook may be used as preparatory training in conjunction with specialist training in female examination such as in intrauterine contraceptive device insertion where knowing the position of the uterus is critical.

Trainers may wish to set a timeframe over which training should be undertaken. The recommendation is that this training period should be no longer than six months.

There is a suggested template for a certificate of competence in Appendix 6 on page 33.

Standards for learning and competency

A competau23

Aid 1 Page

Demonstrates an understanding of how to prepare the environment and equipment for undertaking vaginal, speculum and genital bi-manual examination and specimen collection.
Agree a plan of learning and assessment with trainer/assessor, including timeframes, appointments, reviews and feedback. Comment on, for example: • access to training • training process: sufficient variety of cases • assessment process.
For example, the practitioner will: • prepare or supervise the preparation of the environment, taking into account any specific requirements • ensure environment enables the maintenance of privacy and dignity • prepare the equipment required, eg various sizes of speculum. • ensure all equipment for specimen collection or cytology is present • ensure provision of latex free products • ensure access for women with disabilities such as access to couches and specific positioning requirements.
Describe how the learning strategy is implemented.
Describe how achievement of the learning outcome will be demonstrated to the assessor.

BACK TO CONTENTS

Appendix 2

The learner should keep a logbook of all genital examinations and incidents associated with them, including positive and negative experiences for the learner or woman. This logbook will be used by the learner for learning, study and reflection and for the trainer as an assessment tool.

Name of learner; place of work; contact details; training commencement date
Case number; setting; date seen
Indication for genital examination; history taking; record keeping
Preparation of examination room and woman
Explanation and demonstration of procedure and any tests to be undertaken; how results will be managed
External genitalia examination and findings
Speculum examination and findings

BACK TO CONTENTS

Appendix 3
Skills checklist
for genital examination in women

The assessor should only sign off the appropriate section once the learner is competent. A certificate of competence can be issued by the assessor.

Action	Name and signature of assessor	Date	Name of clinic/ hospital
Part 1			
Has appropriate theoretical knowledge in anatomy, physiology and pathology			
[illegible]			

BACK TO CONTENTS

Activity 5 Self-reflection and learning

Evaluate:
Factors that affected my ability to learn :
Factors that have enabled me to learn :
Areas still to learn more about:
Action plan going forward with time scales and who may be able to help:
Useful resources:

Appendix 6 Sample certificate of competence

Certificate of competence

To be completed by the assessor

I certify that

_____

(learner's name)

Has been assessed at: (name and address of clinic/hospital)

_____

_____

_____

as competent in the following areas:

1 Part 1 Knowledge, attitude, skills: external genitalia examination, speculum examination
Yes/No

2 Part 2 Bimanual examination
Yes/No

Delete as necessary. All learners must complete Part 1.

_____

_____

_____

_____

_____

_____

BACK TO CONTENTS

References

BASHH (2019) Standards for the management of sexually transmitted infections (STIs) , London: BASH. Available at: www.bashh.org/about-bashh/publications/standards-for-the-management-ofstis (accessed 6 September 2023)

Care Quality Commission (2022) GP Mythbusters: Chaperones, London: CQC. Available at: www.cqc.org.uk/guidance-providers/gps/gp-mythbusters (accessed 6 September 2023)

Department of Health (2009) Reference guide to consent for examination or treatment , Second edition, London: DH. Available from: www.gov.uk/government/publications/reference-guide-to-consent-for-examination-or-treatment-second-edition (accessed 6 September 2023)

Department of Health and Social Care (2021) Independent report. Expert Panel on Hymenoplasty. Available at: www.gov.uk/government/publications/expert-panel-on-hymenoplasty (accessed 1 November 2023)

Faculty of Sexual and Reproductive Health (2020) Service Standards for Consultations

Public Health England (2023) Cervical screening: programme and colposcopy management , London: PHE. Available at: www.gov.uk/government/publications/cervical-screening-programme-and-colposcopy-management (accessed 6 September 2023)

Royal College of Nursing (2023) Female genital mutilation: An RCN resource for nursing and midwifery practice , London: RCN. Available at: www.rcn.org.uk/publications (accessed 6 September 2023)

Sexual Offences (Northern Ireland) Order 2008. Available at: www.legislation.gov.uk/nisi/2008/1769/contents (accessed 6 September 2023)

Useful websites

British Association for Sexual Health and HIV
www.bashh.org

British Society for Colposcopy and Cervical Pathology (BSCCP)
www.bsccp.org.uk

Department of Health
www.gov.uk/government/organisations/department-of-health-and-social-care

Faculty of Sexual and Reproductive Healthcare of the Royal College of Obstetricians and Gynaecologists
www.fsrh.org

Institute of Psychosexual Medicine
www.ipm.org.uk

NHS Cervical screening service
www.gov.uk/guidance/cervical-screeningprogramme-overview

Royal College of Obstetricians and Gynaecologists
www.rcog.org.uk

Royal College of Nursing

Modern slavery information

Further reading

British Association for Sexual health and HIV (2023) UK national guideline for consultations requiring sexual history taking

Publication

This is an RCN practice guidance. Practice guidance are evidence-based consensus documents, used to guide decisions about appropriate care of an individual, family or population in a specific context.

Description

This latest edition provides standards and sample assessment tools for training in genital examination in women for registered nurses working in sexual and reproductive health settings, and related health and social care settings.

Publication date: November 2023 Review date: November 2026

The Nine Quality Standards

This publication has met the nine quality standards of the quality framework for RCN professional publications. For more information, or to request further details on how the nine quality standards have been met in relation to this particular professional publication, please contact publications.feedback@rcn.org.uk

BACK TO CONTENTS

The RCN represents nurses and nursing, promotes excellence in practice and shapes health policies

RCN Direct
www.rcn.org.uk/direct
0345 772 6100

Published by the Royal College of Nursing
20 Cavendish Square
London
W1G 0RN

020 7409 3333
www.rcn.org.uk

November 2023
Review date: November 2026
Publication code: 011 162

Royal College of Nursing